

Framingham Pediatrics

Routine Immunization and Screening Schedule

Newborn	HepB #1
2 Week	
1 Month	
2 Month	<u>Vaxelis</u> (DTaP #1, HepB #2, Hib #1, IPV #1) PCV20 #1, Rotavirus #1
4 Month	<u>Vaxelis</u> (DTaP #2, HepB #3, Hib #2, IPV #2) PCV20 #2, Rotavirus #2
6 Month	<u>Vaxelis</u> (DTaP #3, HepB #4, Hib #3, IPV #3) PCV20 #3, Rotavirus #3 (The 6-month visit must be <u>on or after</u> 6 months of age)
9 Month	
12 Month	MMR #1, VARIVAX #1, PCV20 #4 CBC & Lead, SPOT vision screening (The 12-month visit must be <u>AFTER</u> the first birthday)
15 Month	Hib #4, Hep A #1
18 Month	DTaP #4
24 Month	Hep A #2 CBC & Lead, SPOT vision screening (The 24-month visit must be <u>AFTER</u> the second birthday)
2 ½ Year	At the discretion of the physician
3 Year	Hemoglobin & Lead Begin yearly vision screening and BP check
4 Year	Hearing Screening
5 Year	<u>Kinrix</u> (DTaP #5, IPV #5), <u>Proquad</u> (MMR #2, VARIVAX #2)

Yearly physical exams are recommended after 5 years of age.

9 Year	CBC, Cholesterol screening
11 year	Tdap, HPV #1, MCV4 #1 HPV #2 given 6 months later or following year
16 year	MCV4 #2
17 year	CBC, Cholesterol screening

Entrance to 7th Grade requires completion of the Hepatitis B Series, Tdap Booster, documentation of chicken pox disease or Varivax, and 2 MMR vaccines.

MCV4 Booster - Given at age **16** if primary dose is administered at age 11-12yrs, Age 16-18yrs if primary dose is administered at 13-15yrs. **Booster not required if primary dose is administered after 16yrs.**

HPV vaccine if started on or after age 15: 3 dose series (0, 1-2 months, 6 months)

Minimal interval between dose 1 and 2 is 4 weeks, dose 2 and 3 is 12 weeks and dose 1 and 3 is 5 months

The earliest age for 1st dose of **Vaxelis** (DTaP, Hep B, IPV Hib) and PCV20 is 6 weeks old.

1st dose of **Rotavirus** vaccine must be given prior to 12 weeks of age, last dose prior to 32 weeks.

Minimal interval between vaccines **Vaxelis** (DTaP, Hep B, IPV, Hib) and PCV20 is 4 weeks.

Hepatitis B Vaccine for older patients requires minimum 4-week interval between 1st and 2nd doses with 3rd dose given 6-12 months later (minimum intervals of 4 months between 1st and 3rd doses and 2 months between 2nd and 3rd doses).