

Scaphoid Fracture/ Occult Scaphoid Injury



What is a scaphoid injury?

The scaphoid is one of the bones of the wrist, located just below the thumb. Injuries to the scaphoid are some of the most common injuries in children.

A sign that your child has injured their scaphoid is pain right above the scaphoid bone. We sometimes call this "snuffbox tenderness."

This injury occurs most often with what we call a FOOSH (fall onto an out-stretched hand). FOOSH injuries can happen from falls off a scooter, skates or monkey bars, as well as direct hits in sports like football, hockey or lacrosse.

How is this injury treated?

Scaphoid fractures can sometimes be seen on an x-ray, but they are often hard to see, especially during the first two weeks after injury. If there is a break in the bone that cannot be seen on the x-ray, it is called an occult injury.

The scaphoid does not get a large supply of blood, so it can be slow to heal, which can cause complications. Your child may need a cast on their wrist, even if the x-rays appear normal at first.

If your child's provider thinks your child may have a scaphoid fracture but the x-rays are normal, we may order a CT scan or MRI. These tests help us look for other signs that the bone was injured.

If we see a clear fracture, we will determine whether the fracture is displaced, which means that parts of the bone are separated. We usually use a cast for non-displaced fractures and observe them closely. If the fracture is displaced, your child may need surgery, which can give the bone the best chance to heal. Sometimes even nondisplaced scaphoid fractures are treated with surgery.

Will my child be in pain?

Soreness is usually at its worst in the first few days through the first week. Pain from soreness can be treated with acetaminophen (Tylenol®) or ibuprofen (Advil®) as needed. **Always talk with your provider about allergies your child may have before giving over-the-counter medication.**

We may prescribe a small amount of prescription pain medication after a surgery if we feel it is needed.



Swelling in the fingers is common. Have your child keep their arm and hand lifted or resting above their heart.

Can my child be active?


The cast provides some protection, but a blow to the arm could move the fracture out of place or make the injury unstable.

Your child should not participate in activities that put them at risk of falling or a direct hit to the arm. This includes activities like:

- playing on playground structures (i.e. jungle gyms or swing sets)
- contact sports like basketball, hockey or soccer
- horseback riding, ice skating or skiing



Scaphoid fracture



Scaphoid fracture repair

How long will my child be out of sports?

We will assess your child and make recommendations based on how their fracture looks and the potential risks of the sport they play.

Your child probably will not be able to play contact sports or do playground activities for six to 12 weeks, including some recovery time after their cast comes off. If they have a displaced fracture or need surgery, they may need to remain out of these activities for longer.

